

RESEARCH PAPER

## Risk factors of low birth weight in Basra city

<sup>1</sup>Zainab Hussein Al mussa

<sup>2</sup>Narjis A.H Ajeel ,

1 MD, Basrah Health Directorate

2 Dept of Family and Community Medicine, College of Medicine, University of Basrah

Received: 18.04.2021

Accepted: 03.07.2021

### Abstract

**Background** Ready-to-use supplementary foods are high-energy, lipid-based provide energy, protein, fat, vitamins and minerals to treat acute malnutrition in children aged 6 -59 months.

**Aim** To evaluate the effect of ready-to-use supplement foods on the outpatient management of children with acute malnutrition.

**Methods** A prospective appropriate study was carried out on children with acute malnutrition who were referred to the nutritional rehabilitation center at Basra Teaching Hospital; were received ready-to- use supplementary foods at a quantity sufficient to meet their nutrient requirements for full catch- up growth and followed at two subsequent visits

**Results** Moderate wasting and underweight recorded in (66.7 %) and severe wasting in (33.33%) of the patients. Mean weight gain at the first and second follow-up visit was (5.78 ±2.43) and (6.52 ±2.75) g/kg/day respectively. There was a significant improvement in the weight for height Z score at the first and second follow-up visits after the administration of ready-to-use supplementary foods (P value < 0.05). Approximately 32% of the children aged 12-18-month experienced moderate weight gain. Bottle feeding with complementary feeding was reported in 24.19% of the children with moderate weight gain, and 9.52% and 34.68% of the children of illiterate parents and unemployed fathers respectively, showed moderate weight gain. Children belong to families with low income group accounted for 37.09% of those with moderate weight gain. Family income was the only variable that depend on the weight gain results (P value < 0.05). **Conclusion:** Ready-to-use supplementary food is significantly effective for outpatient management of acute malnutrition.

**Keywords:** Acute Malnutrition, Ready-to-Use Supplementary Food, Outpatients

### Corresponding to:

Dr. Zainab Hussein Al Mussa, MD, Basrah Health Directorate

✉ [zalmussazyahoom@gmail.com](mailto:zalmussazyahoom@gmail.com)

### Introduction

Low birth weight (LBW) (a birth weight of less than 2500 grams), is one of the best indicators of health risk and the best predictor of mortality and morbidity in the neonatal, post neonatal and childhood periods. <sup>(1)</sup>

LBW is one of the leading causes of infant mortality and morbidity, and those infants are 40 times more likely to die in their first month of their life than normal birth weight infants. They are also twice as likely as other infants to exhibit health problems and serious developmental delays during childhood <sup>(2)</sup>

It's also associated with immediate and late sequelae. Immediate complications include: hypoxia, ischemia, sensorineural injury, anemia, respiratory failure, and microcephaly. While late

complications include: epilepsy, hydrocephalus, osteopenia, fractures, hearing and visual impairment<sup>(3)</sup>

very low birth weight (VLBW) is a birth weight less than 1,500 gm at birth and predominately immature<sup>(3)</sup>, is an accurate predictor of infant mortality rate (Relative risk of 93). VLBW infants in USA account for over 50% of neonatal deaths and 50% of handicapped infants; their survival is directly related to birth weight.<sup>(1)</sup>

In 2015 estimate that 20.5 million (uncertainty range 17.4–24.0) livebirths had a birthweight of less than 2500 g. Most (91%) were in low-income and middle-income countries.<sup>(4)</sup>

In Latin America and the Caribbean and Western Europe, there was no apparent change in low birthweight prevalence between 2000 and 2015. Similarly, reported data from 57 mostly high-income countries with relatively low baseline suggests almost no change in LBW prevalence. For the remaining countries, a 17% reduction in LBW prevalence was estimated over the years 2000–15, most notably in the countries with the highest prevalence.<sup>(4)</sup>

According to UNICEF the percentage of low birth weight infants in Middle East and South Africa in 2000 was 15%.<sup>(2)</sup> Central and South America have, on average, much lower rates (10%), while in the Caribbean the level is 14%.<sup>(4)</sup>

In developing countries, approximately 70% of LBW infant have IUGR.<sup>(2)</sup> In Iraq Up to 1990, the 5% incidence of LBW was similar to industrialized countries. Since then, a marked increase to 22% in 1995 was reported.<sup>(2)</sup> In 2017, it was reported to be 20.2%.<sup>(5)</sup>

**The aims of the study:** The present study was carried out to study the association between birth weight and selected risk factors in Basrah governorate with special reference to socio-

demographic factors, present and past obstetric history, and maternal medical history.

## Methods

This is a hospital- based cross sectional study which was designed to study the association between LBW and selected risk factors. The study involved infants delivered and /or admitted to AL-Basrah-Maternal and Child Hospital during the period from the first of May to the 31st. of October 2007 on three selected days / week during the whole study period. A total of 400 infants were initially included, however 38 twins were excluded ( because the study for birth weight of singleton births only )leaving 362 singleton births to be included in the study.

The data were collected using a special questionnaire form designed (by researchers ) for the purpose of the study. The questionnaire was designed to provide information about the following aspects: socio-demographic characteristics of the parents (age, education, occupation), birth weight, birth order, preceding birth interval, history of twin, history of recent pregnancy complications, past medical history, past obstetric history and history of Antenatal care..

The questionnaire form was filled by one of the authors through direct interview of the mothers. Before each interview, the purpose of the study was clearly explained to the mothers and their verbal consent was obtained. No objection was faced by any woman giving a response rate of 100%. Each interview took about fifteen minutes. The birth weight for included infants were either measured by one of the authors for recent births delivered in the hospital by using the same weighing scale ( uniscale ) or recorded as

reported by mothers for infants admitted to the hospital.

**Statistical analysis:** The data were analyzed using SPSS (Statistical Package for Social Sciences) program version 15. Chi squared test was used to find out the significant associations between different variables. A P- value of < 0.05 was the criterion of statistical significance.

## Results

### Characteristics of the studied infants

Table 1 showed the characteristics of the studied infants. Out of the 362 infants included in the study, 215(59.4%) were males and 147(40.6%) were females. One-hundred and forty-five (40.1 %) were first born, while 79 (21.8 %) were fifth or later born. Nearly half (48.1%) were born with LBW, the mean birth weight of the studied infants was 2.49 kgm ± 0.49 SD.

TABLE (1) CHARACTERISTICS OF THE STUDIED INFANTS		
Variable	No. ( 362)	%
<b>Sex</b>		
Male	215	59.4
Female	147	40.6
<b>Birth order</b>		
1 <sup>st</sup>	145	40.1
2-4	138	38.1
≥5	79	21.8
<b>Birth weight (kgm)</b>		
<2.5	174	48.1
≥2.5	188	51.9

Mean± SD	2.49±0.49	
Total	362	100.0

### Risk factors of LBW

#### Maternal age

Table 2 shows the association between birth weight and maternal age at time of birth. The highest percentage of LBW was for births to mothers in the age group 30- 39 years. However, the association between maternal age and birth weight was statistically not significant (P> 0.05).

Table (2) The relationship between birth weight and maternal age						
Maternal age (years)	Birth weight (kgm)				Total	
	< 2.5		≥2.5			
	No.	%	No.	%	No.	%
< 20	38	47.5	42	52.5	80	100.0
20-29	71	42.8	95	57.2	166	100.0
30-39	58	58.6	41	41.4	99	100.0
≥40	7	41.2	10	58.8	17	100.0
Total	174	48.1	188	51.9	362	100.0
$\chi^2=6.587$ df=3 p value= 0.086						

#### Maternal education and employment

A significant association between maternal education and birth weight was found (p < 0.01). A clear consistent decline in the incidence of LBW was observed with the highest incidence of LBW was found among births to low educated or uneducated mothers (finished their primary education or less including illiterate), while the lowest incidence was among births for mothers who had ≥10 years of education (Table 3). On the other hand, no significant association was found

between maternal employment and low birth weight (P value > 0.05)

### **Complications during pregnancy**

Table (4) shows the relationship between history of complications during pregnancy and birth weight. It's evident that a higher percentage of mothers who had complications during pregnancy (hypertension, infection, bleeding) gave birth to babies with LBW (68.9%) compared to those who had no complication (24.3%) the association was statistically highly significant ( $p < 0.001$ ).

### **Previous history of LBW**

Table (5) shows that mothers who had previous history of having LBW infants were more likely to report a LBW for the current birth (67.7 %) compared to those with no such a history (45.2 %). The association between previous history of LBW and subsequent risk of LBW was statistically significant ( $P < 0.01$ ).

### **Gestational age**

Table (6) shows that the percentage of LBW was markedly higher among preterm births (gestational age < 37 weeks) than that among full term births (gestational age  $\geq$  37 weeks), 72.6 % and 34.0 % respectively. The association between gestational age and birth weight was statistically highly significant ( $p < 0.001$ ).

### **Preceding birth interval**

The relation between preceding birth intervals and birth weight is shown in Table (7). The highest incidence of LBW (73.5%) was for infants born after a very short preceding birth interval ( $\leq$  6 months). The incidence gradually decreased to its lowest level when the preceding

birth interval was in the range between 13-24 month (37.9%). The incidence of LBW then increased again when the preceding birth interval was > 24 months. Statistically there is significant association between preceding birth intervals and birth weight ( $P < 0.05$ )

### **Logistic regression Analysis**

When the logistic regression analysis was carried out, it was found that only preterm labour and complications during pregnancy were independently and significantly affected the birth weight. Table (8)

## **Discussion**

### **Risk factors**

#### **Maternal age**

The present study showed that the highest percentage of LBW was for births of older mothers. Worldwide some studies reported a higher risk of LBW among older mothers. <sup>(6,7)</sup> While others found that teenage mothers (mothers less than 20 years of age) were more likely to have LBW than mothers in other age groups. <sup>(8,9,10)</sup> There were many factors that contribute to this risk for example, young teenagers often have poor eating habits, poor weight gain, and neglect to take their vitamin supplements, and fail to seek prenatal care early in pregnancy. They may also smoke. As women get older their chance of multiple gestation (twin, triple, etc.) increases, old women are also prone to have hypertension, DM, and cardiac disease, these diseases can complicate her pregnancy and make her more prone to have LBW baby. <sup>(8,11)</sup>

#### **Maternal education and employment**

Regarding maternal education, the present study showed that the highest percentage of LBW in births for mothers with low educational level. This is comparable to the results of a study

carried out in north – west Russia <sup>(6)</sup>, which showed the effect of patient's education on the risk of LBW. This might be explained by the fact that the educated mothers were more aware about their diet during pregnancy, the importance of early and adequate (Antenatal care) ANC and the importance of spacing between pregnancies <sup>(12)</sup>. While some studies found that maternal occupation was one of the risk factors for LBW, others found that maternal occupation did not influence the risk of LBW <sup>(13,14)</sup>. In the present study, the risk of LBW was higher among births for employed mothers. This may be due to the physical stress associated with some occupations for example, mothers who stand for long periods of time or involved in hard work may have chance of increase uterine contraction, premature rupture of membrane and a higher rate of preterm delivery and LBW. <sup>(13)</sup>

#### **Complications during pregnancy**

It was evident in the present study that a higher percentage of mothers who had complications during pregnancy gave birth to babies with LBW compared to those who didn't develop such complication. The association between complications during pregnancy and the risk of LBW was statistically significant and independent on other risk factors. This is expected and it agrees with many other studies. <sup>(15,16,12,17)</sup> Hypertension can constrict the blood vessels in the uterus that supply the fetuses with oxygen and nutrients, when this occurs before term, it can slow the fetus growth, sometimes resulting in low birth weight, hypertension also increases the risk of preterm delivery. <sup>(18)</sup> Infections especially those involving the genitourinary tract, may increase the risk of preterm delivery. <sup>(17)</sup> Infection with certain viruses, including cytomegalovirus, rubella and

chickenpox can also slow fetal growth and cause birth defects. <sup>(19)</sup>

#### **Gestational age**

Many factors affect the duration of gestation and fetal growth, thus birth before 37 weeks of gestation (preterm birth) is a major component of low birth weight. <sup>(14)</sup> The present study showed that a higher percentage of LBW among preterm births, the association between preterm delivery and the risk of LBW was statistically significant and independent on other risk factors. This is in agreement with other studies. <sup>(20, 21)</sup> Preterm delivery leads to LBW because those births had restricted period of intrauterine growth. <sup>(22)</sup>

#### **Previous history of LBW**

Mothers who had previous LBW babies had higher rate of having another LBW baby. <sup>(12, 23)</sup> This may be due to the fact that those mothers may be still exposed to the same risk factors which lead in LBW in the previous pregnancies.

#### **Preceding birth intervals**

A very short preceding birth interval could be a leading cause for LBW, in this study a higher percentage of LBW was found among mothers with very short preceding birth intervals. This was identical to the finding of a study in Nevada in USA, which found that an infant born within 12 months of previous birth was more likely to be of LBW than infants born 24 months after a previous live birth. This can be explained that mother body did not get enough time to recover after last pregnancy and this exposed her to risk of anemia and low pregnancy weight gain. <sup>(5)</sup>

#### **Limitations of the study:**

This study is a cross sectional study based on 362 newly born infants. The study was carried out in AL – Basrah Maternity and child hospital which is the main maternity hospital in Basrah city.

However, the studied births may not be representative of all births in the city.

A recall bias may also be a problem in obtaining information on variables such as birth weight and frequency of ANC visits. The effect of recall bias was minimized by extracting some of the information from the ANC card or from the case sheet.

Another source of bias is the non-response bias, but fortunately in the present study, the response rate was 100% because all women were willing to participate in the study.

## **Conclusion**

The factors which showed a significant independent association with LBW were complications during pregnancy and preterm delivery.

**Conflict of interest:** The authors declare no conflict of interest.

**Financial Support:** None declared.

TABLE (3) THE RELATIONSHIP BETWEEN BIRTH WEIGHT AND MATERNAL EDUCATION & EMPLOYMENT						
Educational level(years)	Birth weight (kgm)				Total	
	<2.5		≥2.5			
	No.	%	No.	%	No.	%
≤6	110	51.5	98	48.5	208	100.0
7-9	45	48.9	48	51.1	93	100.0
≥10	19	31.1	42	68.9	61	100.0
Total	174	48.1	188	51.9	362	100.0
$\chi^2=13.098$ df= 2      p value = 0.01						
Maternal employment	Birth weight (kgm)				Total	
	<2.5		≥2.5			
	No.	%	No.	%	No.	%
Non employed	158	47.0	178	53.0	326	100.0
Employed	16	61.5	10	38.5	26	100.0
Total	174	48.1	188	51.9	362	100.0
$\chi^2=2.037$ df = 1      p value = 0.154						

Table (4) the relationship between complications during pregnancy and birth weigh						
Complications during pregnancy	Birth weight (kgm)				Total	
	< 2.5		≥ 2.5			
	No.	%	No.	%	No.	%
Positive	133	68.9	60	31.1	193	100.0
Negative	41	24.3	128	75.7	169	100.0
Total	174	48.1	188	51.9	362	100.0
$\chi^2=71.965$ df= 1      p value =0.00						



**TABLE (8) THE LOGISTIC REGRESSION ANALYSIS**

Variables in the Equation	OR	95%CI	P
Age	0.975	0.975-1.041	0.445
Parity	1.210	0.999-1.466	0.051
Education	1.054	0.973-1.141	0.198
occupation(1)	0.774	0.238-2.519	0.670
Birth interval	0.995	0.978-1.012	0.547
preterm(1)	28.994	12.436-67.599	0.000
Baby sex(1)	0.841	0.475-1.489	0.552
Antenatal care(1)	0.846	0.382-1.875	0.681
Previous low birth weight (1)	1.818	0.821-4.024	0.141
Present complication during pregnancy (1)	6.634	3.740-11.769	0.000
Constant	0.194		0.120

## References

- 1.WHO. Guidelines on optimal feeding of low birth-weight infants in low-and middle-income countries. World Health Organization, 2011.
- 2-. Nelson text book of pediatrics 19th .obert M. Kliegman, MD ,Professor and ChairDepartment of Pediatrics Medical College of Wisconsin Joseph W. St. Geme III, MD, Nina F. Schor, MD, PhD ,William H. Eilinger Professor and Chair,Nonprofit Healthcare and Educational ,Consultants to Medical Instituti , Philadelphia; WBSaunders,2015.p. 476-479 .
- 3-Jammeh, Abdou, Johanne Sundby, and Siri Vangen. "Maternal and obstetric risk factors for low birth weight and preterm birth in rural Gambia: a hospital-based study of 1579 deliveries." *Open Journal of Obstetrics and Gynecology* 1.03 (2011): 94.
- 4-Blencowe, H., Krusevec, J., de Onis, M., Black, R. E., An, X., Stevens, G. A., ... & Cousens, S. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health* 2019; 7
- 5- HAMZA, Nazhat Jassab. Causes of Neonatal Mortality in Neonatal Care Units AL-Diwaniya Province–Iraq 2013. *Al-Qadisiyah Medical Journal*, 2019, 15.1: 76-86.
- 6-Usynina, A. A., Grjibovski, A. M., Krettek, A., Odland, J. Ø., Kudryavtsev, A. V., & Anda, E. E. (2017). Risk factors for perinatal mortality in Murmansk County, Russia: a registry-based study. *Global health action*, 10(1), 1270536.
7. A-Dekna M,AL-Hadithi TS,A.AL- Diwan JKA, Shaban ALO. Significance of birth weight in viability of the infected newborn in Tripoli Medical Center. *J Fac of Med (Baghdad)* 202; 44(3):19-22.
- 8-Aras, Radha Y. "Is maternal age risk factor for low birth weight?." *Archives of medicine and health sciences* 1.1 (2013): 33.
- 9-da Roza, Daiane Leite, Carla Maria Teixeira de Oliveira, and Edson Zangiacomi Martinez. "Spatio-temporal trends in the risk of adolescent pregnancy in Minas Gerais, Brazil, 2000–2015." *International journal of adolescent medicine and health* 1.ahead-of-print (2018).
10. Feris BG. Epidemiology standardization project (American thoracic society). *J AM Rev Respir Dis* 1978;(1):118-120.
11. GURAV RB , KARTIKEYAN S, JAPE MR.Pilot study on LBW risk factors .[online].2003[cited July; 2003] ;Available from:  
[URL:http://www.bhj.org/journal/20034503july/owbirth413.htm](http://www.bhj.org/journal/20034503july/owbirth413.htm)
- 12.Majoko F, Nystrom L, Munjanja S, Lindmark G. Usefulness of risk scoring at booking for ANC in predicting adverse pregnancy Out come in a rural African setting . *J Obstet Gynaecol* 2002 ; 22(6):604-9.
- 13-Williams Obstetrics: 23rd Edition 2013 ,by F. Cunningham (Author), Kenneth Leveno (Author), Steven Bloom (Author), John Hauth (Author), Dwight Rouse (Author), Catherine Spong (Author), page 601,959.
- 14.Rondo PH, Ferreira RF, Nogueira F, Ribeiro MC, Lobert H, Artes R. Maternal psychological stress and distress as predictors of LBW , prematurity and intrauterine growth retardation, Brazil. *Eur J Clin Nutr* 2003 ; 57(2): 266-72.(midline)
- 15- Figueiredo, A. C. M. G., Gomes-Filho, I. S., Batista, J. E. T., Orrico, G. S., Porto, E. C. L., Cruz Pimenta, R. M., ... & Pereira, M. G. (2019). Maternal anemia and birth weight: A prospective cohort study. *PloS one*, 14(3), e0212817.

- 16- Lumbiganon, P., Laopaiboon, M., Gülmezoglu, A. M., Souza, J. P., Taneepanichskul, S., Ruyan, P., ... & Villar, J. (2010). Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08. *The Lancet*, 375(9713), 490-499.
- 17-Koc, E., Demirel, N., Bas, A. Y., Ulubas Isik, D., Hirfanoglu, I. M., Tunc, T., ... & Hakan, N. (2019). Early neonatal outcomes of very-low-birth-weight infants in Turkey: A prospective multicenter study of the Turkish Neonatal Society. *Plos one*, 14(12), e0226679.
- 18-Harper, L. M., Jauk, V., Longo, S., Biggio, J. R., Szychowski, J. M., & Tita, A. T. (2020). Early gestational diabetes screening in obese women: a randomized controlled trial. *American journal of obstetrics and gynecology*, 222(5), 495-e1.
- 19- FAREWELL, Vern; JOHNSON, Anthony. The origins of Austin Bradford Hill's classic textbook of medical statistics. *Journal of the*

- Royal Society of Medicine*, 2012, 105.11: 483-489.
- 20-Cruz, H. O., Plascencia, J. L., Nakashimada, A. I., & Espinoza, J. M. G. (2007). Causes of preterm delivery among adolescent mothers. *Ginecologia y obstetricia de Mexico*, 75(01), 17-
21. Al.Satar AA. Study of The Morbidity and Mortality among LBW infants in AL -Basrah Maternal and Child Hospital. Diploma report, collage of medicine ,university of Basrah2001.
- 22-Mayes' Midwifery: A Textbook for Midwives 14th Edition Editor: Sue Macdonald Book ISBN: 9780702045967 Imprint: Baillière Tindall Published Date: 19th April 2011. 852–857.
23. AL.Rubaiy M. Risk factors and Labour outcome in parous women with Pre-Eclampsia. *Medical Journal of Basrah university* 2006; 24(1,2):25-26