

LIFESTYLE DETERMINANTS OF HIGH- DENSITY LIPOPROTEIN CHOLESTEROL (HDL-C) IN YOUNG ADULTS

Jasim N. Al-Asadi, Omran S. Habib & Lamia M Al-Naama

ABSTRACT

High-density lipoprotein cholesterol is a potent predictor of coronary heart disease. Genetic as well as environmental factors including lifestyle factors play a role as determinants of its level in the blood. To examine the effects of certain lifestyle factors on serum level of high density lipoprotein cholesterol in young adult people. Three-hundred thirty-eight young adult subjects of both sexes aged 20-44 years asymptomatic for cardiovascular diseases were interviewed according to special questionnaire including information on lifestyle habits. Physical examination was done, height, body weight, and blood pressure measurements were performed. Blood analysis to determine the blood level of high density lipoprotein cholesterol was done after 12 hours fasting. Smoking and obesity were the most significant risk factors associated with a decreased level of high density lipoprotein cholesterol. The level of HDL-C was 50.9 ± 12.1 mg/dl in smokers compared with 58.3 ± 12.8 mg/dl in non-smokers. Its level was 49.8 ± 9.8 mg/dl in obese individuals compared to 58.4 ± 12.8 mg/dl in normal body weight subjects. Physical activity was not significantly associated with low level of HDL-C by univariate analysis, but it was found to be significantly associated with its level by the multiple regression analysis. High-density lipoprotein cholesterol level was a function of many factors, some of them were lifestyle related such as smoking, physical activity, and obesity. Therefore, efforts to encourage more physical activity, quitting smoking, consuming low fat diet, and keeping ideal body weight are recommended.

INTRODUCTION

Evidence from epidemiological and clinical trials had supported an inverse relationship between HDL-C and risk of coronary heart disease (CHD).^[1,2] For every 1mg/dl increase in serum HDL-C, there appears to be a corresponding 2% to 3% decrease in CHD risk and 4% to 5% decrease in cardiovascular diseases mortality.^[1] Approximately one quarter to one third of patients with pre-existing coronary disease and desirable total cholesterol [less than 5.2 mmol/L] have low levels of HDL-C [less than 1 mmol/L] as the primary abnormality.^[3] The protective effects of HDL-C are multifactorial and many possibilities are suggested; HDL-C may prevent oxidation of LDL-C, so it protects against excess lipid accumulation in the blood vessel wall.^[4] It provides a reverse cholesterol transport from the tissues to the liver for metabolic conversion and excretion.^[5] It has also been hypothesized that cholesterol efflux from atherosclerotic lesion is promoted by HDL-C, possibly through a receptor related mechanism.^[6] In addition to constitutional determinants such as age, sex, ethnicity and genetic factors, other variables such as dietary fat, obesity and other lifestyle factors including

physical activity, alcohol consumption and smoking habit are known to affect serum HDL-C level.^[7-11] There are many studies worldwide which studied the influence of lifestyle on serum lipids (including HDL-C),^[9-11] but studies which examined the relation between lifestyle factors and HDL-C serum levels in young adults asymptomatic for coronary heart diseases are scarce in Iraq.^[12] Therefore this cross-sectional study was conducted in Basrah (in 2003) to examine the effect of certain lifestyle factors on serum HDL-C level.

SUBJECTS AND METHODS

Subjects: The study population consisted of 338 young adult individuals of both sexes (aged 20-44 years) apparently healthy, with no history of cardiovascular diseases, who were selected randomly from three colleges from the University of Basrah and from Iraqi divers.

Methods

Interviewing was performed according to a special questionnaire form which covers the following aspects; Socio-demographic characteristics, medical history, aspects of dietary habits, alcohol consumption, physical

activity, and smoking. Smoking habits: non-smokers defined by those who never smoke, Ex-smokers included those who stopped smoking before more than 3 months, while Current smokers were those who smoke regularly. Diet was classified on the type of food that individuals consumed, those who consumed a well known fatty diet such as butter, cream, cheese, solid fat---etc, on most days of the week were considered to be on fatty diet. Physical activity was measured by using a combined index of leisure time physical activity which was calculated from the product of intensity, estimated duration of exercise and monthly frequency using the method reported by Raitakan et al^[13]. The range of index is 1-225. Subjects with index of equal or higher than 85 were considered as a constantly active, which nearly equals to an intensive physical activity for more than 2 hours/week. Subjects with an index value of less than or equals to 15 were considered as constantly sedentary which nearly equals to 1 hour of light aerobic activity/ week. Those who have an index of 16-84 were considered moderately active.^[13] Height and weight were measured, and the body mass index (BMI) was calculated using the Quetlet index [BMI= weight in Kg/height in m²]^[14] Non obese: BMI <25kg/m², overweight: BMI 25-29.9 kg/ m², and obese if BMI was ≥30 kg/ m².^[15] Clinical heart examination and pulse rate measurements were done to detect any heart abnormality. Biochemical measurements regarding serum HDL-C estimation was performed after 12 hours fasting. Serum HDL measurement was done by the bioMerieux (Kit No. 61 231) after precipitation and then estimating the cholesterol content depending on the enzymatic method of Bruestien et al.^[16] Low level of HDL-C was considered if the level was < 40 mg / dl.^[17]

Statistical analysis

The statistical analysis was made by the use of the Statistical Package for Social Science (SPSS) version 11. Analysis of variance and t-test were used when appropriate. Multiple regression analysis was used to determine whether observed differences in outcomes remained while controlling for potentially confounding variables. A P-value of <0.05 was considered significant.

RESULTS

Table-1 demonstrates selected socio-demographic characteristics of the study population. The majority were in their early twenties. While those who were above 40 years of age formed only 3% of the total studied population. Regarding sex composition, the majority were men. 63.5% were with secondary school level of education, and 13.1% were with basic university or more level of education.

Table 1. Socio-demographic characteristics of the study population

Character	No.	%
Age (years)		
20 – 24	209	61.7
25 – 29	59	17.5
30 – 34	32	9.5
35 – 39	28	8.3
40 - 44	10	3.0
Sex		
Men	279	82.5
Women	59	17.5
Residence		
Urban	251	74.3
Rural	87	25.7
Education		
Illiterate	8	2.4
Primary & Intermediate	71	21.0
Secondary	215	63.5
Basic university and above	44	13.1
Marital status		
Married	87	25.7
Single	251	74.3
Total	338	100

Tabl-2 shows the frequency of the lifestyle factors among the studied subjects. The fatty diet consumption was the most prevalent lifestyle among the study population. It was prevalent in 34.3% of them, followed by smoking which was prevalent in 24.3%. Obese and inactive subjects formed 9.8% and 8.3% of the total studied population respectively.

Reliable information concerning alcohol intake was difficult to obtain from interviewees, therefore, the results were undependable. Only 3 individuals (0.9%) mentioned they were on alcohol-intake, so no further analysis was carried out regarding this variable.

Table 2. Frequency of the lifestyle determinants among the study population

Determinant	No.	%
Smoking		
Non-Smokers	245	72.5
Ex-smokers	11	3.2
Smokers	82	24.3
Physical activity		
Active	146	43.2
Moderately active	164	48.5
Inactive	28	8.3
Diet		
Non-fatty diet	222	65.7
Fatty diet	116	34.3
BMI		
Normal < 25	232	68.5
Overweight 25-29.9	73	21.6
Obese ≥30	33	9.8
Alcohol		
No Alcohol-Intake	335	99.1
Alcohol-Intake	3	0.9
Total	338	100

As shown in Table-3, HDL-C level was inversely associated with smoking and body mass index with a highly significant difference. HDL-C serum level was 50.9±12.1 mg/dl in smokers in comparison with 58.3±12.8 mg/dl in non-smokers, and its level was 58.4±12.8 mg/dl in normal weight individuals compared to 49.8±9.8 mg/dl in obese persons. While HDL-C level was lower in fatty diet consumers (55.2±12.5 mg/dl) and sedentary individuals (55.7±10.6 mg/dl) in comparison with non-fatty diet consumers (57.03±13.3 mg/dl) and active individuals (57.1±13.8 mg/dl) respectively, but the differences were not significant.

Table 3. HDL-C level according to the studied lifestyle factors

Determinant	Mean ± SD (mg/dl)	P-value
Smoking		
Non-smokers	58.3 ± 12.8	<0.001
Ex-smokers	55.3 ± 13.6	
Smokers	50.9 ± 12.1	
Physical activity		
Active	57.1 ± 13.8	NS
Moderately active	56.2 ± 13.01	
Inactive	55.7 ± 10.6	
Diet		
Non-fatty diet	57.03 ± 13.3	NS
Fatty diet	55.2 ± 12.5	
BMI (Kg/m²)		
Normal < 25	58.4 ± 12.8	<0.001
Overweight 25-29.9	53.4 ± 13.5	
Obese ≥30	49.8 ± 9.8	

To investigate the independent effects of selected risk factors on HDL-C level, a stepwise linear multiple regression analysis was performed (Table-4) the examined risk factors were the lifestyle determinants (i.e. smoking, physical activity, diet, and body mass index) in addition to age and sex. Body mass index, smoking and physical activity in addition to sex (females showed higher HDL-C level), and age were noticed to have highly significant effects on HDL-C concentration. BMI appears to be the strongest variable that explained 6.5% of the variability in HDL-C concentration, together with the other variables, they explained 15.9% of the variation in HDL-C concentration and about 84% of the variation is still left unaccounted for.

Table 4. Multiple regression analysis

Variable	Beta	R2	P-value
BMI	- 0.160	0.065	0.000
Smoking	- 0.181	0.104	0.000
Age	- 0.155	0.122	0.006
Sex	0.191	0.138	0.007
Physical activity	0.163	0.159	0.002

DISCUSSION

This study has shown different contributions to the risk of low level of HDL-C from the major lifestyle factors. Smoking and HDL-C were inversely related, a result that had been reported in young people.^[18,19] This confirms the finding that smoking is associated with an unfavorable lipoprotein profile at whatever age; smoking is initiated during adolescence.^[20] In accordance with the findings of other studies,^[21,22] BMI (an indicator of lifestyle habits such as physical inactivity and diet) was inversely associated with HDL-C, a result that was noted not only by univariate analysis, but also replicated by multiple regression analysis. The finding that physical activity was not significantly related to HDL-C which was noted in univariate analysis dropped out in the stepwise multiple regression analysis. This finding is consistent with that of Savig and Goldbourt.^[11] Diet was the only lifestyle factor which was not found to contribute greatly to a significant HDL-C reduction. Although diet was suggested to be

the major environmental factor that modulates lipid and lipoprotein,^[23] but the relation between fatty diet consumption and HDL-C level in this study was not found to be significant. However, its level was lower in fatty diet consumers in comparison with non-fatty diet consumers. Probably the relative proportion of carbohydrate to fat in the diet is more important factor than estimating the effect of dietary fat content alone (higher ratio is correlated with low HDL-C).^[24] Also, favorable but not significant increase in HDL-C concentration was observed with modified lower carbohydrate diet.^[25] Dietary habit was not quantitatively measured, in this sense, inaccuracy was unavoidable, and thus may have failed to detect a real association between fatty diet and HDL-C level. As shown in the multiple regression analysis only 15.9% of the variation in HDL-C concentration was related to lifestyle factors, sex, and age. While about 84% of the variation in HDL-C left unaccounted for. This may explain the important effect of other environmental and genetic factors on HDL-C serum level. Genetic factors are known to play a major role as determinants of HDL-C, with estimate of heredity from about 40% to 60%.^[26,27]

In conclusion, HDL-C level was primarily a function of many factors, some of them are lifestyle related such as smoking, physical activity and obesity. The results of this study clearly demonstrated the need to initiate preventive efforts early in life to encourage more physical activity, quitting smoking, keeping ideal body weight, and consuming healthy diet.

REFERENCES

1. Sacco RL, Benson RT, Kargman DE, Boden-Albala B, Tuck C, Lin IF, et al. HDL-C and Ischemic stroke in the elderly: The Northern Manhattan Stroke Study JAMA 2001; 285: 2729-2735.
2. Aztalas BF, Schaefer EJ. HDL sub populations in pathological conditions. Amer J Cardiology 2003; 99 (suppl.): 12E-17E.
3. Rubins HB, Robins SJ, Collins D, Iranmanesh A, Timothy J, Mann D, et al. Distribution of lipids in 8500 men with coronary artery disease. Am J Cardiology 1995; 75: 1202-1205.
4. Barter P. Is high-density lipoprotein the protector of the cardiovascular system? European Heart Journal supplements 2004; 6 (supplement A): A 19-A 22.
5. Miller M. Raising an isolated low HDL-C level: Why, how, and when? Cleveland Clinic Journal of Medicine 2003; 70(6): 553-560.
6. Fuster V, Gotto AM, Libby P, Loscalzo J, McGill HC. Pathogenesis of coronary disease: The biologic role of risk factors. 27th Bethesda conference Task Force 1. J Am Coll Cardiology 1996; 27: 964-976.
7. Namekata T, Moore DE, Suzuki K, Mori M, Knopp RH, Marcovina SM, et al. Biological and lifestyle factors and lipid and lipoprotein levels among Japanese Americans in Seattle and Japanese men in Japan. International Journal of Epidemiology 1997; 26(6): 1203-1213.
8. De Oliveira Silva ER, Kong M, Han Z, Starr C, Kass EM, Juo SH, et al. Metabolic and genetic determinants of DL metabolism and hepatic lipase activity in normolipidemic females. Journal of Lipid Research 1999; 40: 1211-1221.
9. Ellison RC, Zhang Y, Qureshi MM, Knox S, Arnett DK, Province MA. Lifestyle determinants of high-density lipoprotein cholesterol: The National Heart, Lung, and Blood Institute Family Heart Study. Am Heart Journal 2004; 147(3): 529-535.
10. Wilsgaard T, Arnesen E. Change in serum lipids and body mass index by age, sex, and smoking status: The Tromsø study 1986-1995. Annals of Epidemiology 2004; 14(4): 265-273.
11. Sagiv M, Goldbourt U. Influence of physical work on high-density lipoprotein cholesterol- Implications for the risk of coronary heart disease. International Journal of Sports Medicine 1994; 15(5): 261-266
12. Ajlaan SK. Lipid profile and physical activity. J Bahrain Medical Society 2004; 16(4): 186-191.
13. Raitakan OT, Porkka KVK, Taimela S, Telama R, Rasanen L, Viikari JS. Effects of persistent physical activity and inactivity on coronary risk factors in children and young adults: The Cardiovascular Risk in Young Finns Study. Amer J Epidemiol 1994; 140(3): 195-205.
14. Frier BM, Truswell AS, Shepherd J, et al. Nutritional factors in health and disease. In: Haslett C, Chillver ER, Hunter TA & Boon NA (Eds): Davidson's principles and practice of Medicine. 18th ed. London. Churchill living stone. 2001.
15. WHO: Obesity, preventing and managing the global epidemic. Technical reports series 894, WHO, Geneva, Switzerland, 2000.
16. Burstein M, Scholnick HR, Morfin R. Rapid method for the isolation of lipoproteins from human serum by precipitation with polyanions. J Lipid Res 1970; 11:583-595.
17. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report on the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). JAMA 2001; 285(19): 2486-2497.
18. Miller ER, Appel LJ, Jiang L, Risby TH. Association between cigarette smoking and lipid peroxidation in a controlled feeding study. Circulation 1997; 96: 1097-1101.
19. Sharma SB, Dwivedi S, Prabhu KM, Singh G, Kumar N, Lal MK. Coronary risk variables in young asymptomatic smokers. Indian J Med Res 2005; 122: 205-210.

20. Clarke WR, Srinivasan SR, Shear CL, Hunter SM, Croft JB, Webber LS, Berenson GS. Cigarette smoking initiation and longitudinal changes in serum lipids and lipoproteins in early adulthood; The Bogalusa Heart study. *Amer J Epidemiol* 1986; 124: 207-219.
21. Plourde G. Impact of obesity on glucose and lipid profile in adolescents at different age groups in relation to adulthood. *BMC Family Practice* 200; 3: 18-31.
22. Assmann G, Schulte H. Relation of HDL-C and triglycerides to incidence of atherosclerotic CAD (The PROCAM experience): A prospective cardiovascular Munster study. *Am J Cardiol* 1992; 70(7): 733-737.
23. Schafer EJ. Lipoproteins, nutrition, and heart disease. *Amer J Clin Nutr* 2002; 75:161-212.
24. Gordon DJ, Witztum JL, Hunninghake D, Gates S, Glueck CJ. Habitual physical activity and HDL-C in men with primary hypercholesterolemia: The Lipid research Clinics Coronary Primary Prevention Trial. *Circulation* 1983; 67(3): 512-520.
25. Aude YW, Agatston AS, Lopez-Jimenez F, Lieberman EH, Almon M, Hansen M, et al. The national cholesterol education program Diet vs a Diet Lower in Carbohydrate and Higher in Protein and Monosaturated Fat. *Arch Intern Med* 2004; 164: 2141-2146.
26. Cohen JC, Wang Z, Grundy SM, et al. Variation at the hepatic lipase and apolipoprotein A1/CIII/AIV loci is a major cause of genetically determined variation in plasma HDL-Cholesterol levels. *J Clin Invest* 1994; 94: 2377-2384.
27. Inazu A, Nishimura Y, Terada Y, Mabuchi H. Effects of hepatic lipase gene promoter nucleotide variations on serum HDL-Cholesterol concentration in the general Japanese population. *J Hum Genet* 2001; 46(4): 172-177.